

Topic: Logarithm

- 1 Given that $u = \log_3 z$, find, in terms of u ,
 - (a) $\log_3 9z$, [1]
 - (b) $\log_3 \left(\frac{z}{27} \right)$, [1]
 - (c) $\log_z 27$. [2]

- 2 Solve
 - (a) $\log_2 (3x - 5) + 3 = \log_2 (4x + 5)$, [3]
 - (b) $2\log_3 y - \log_y 27 = 1$. [5]

- 3 Solve $2\log_7 p = 3 + \log_p 49$. [5]

- 4 Express $2\log_5 x - \log_5 (x - 6) = 1$ as a quadratic equation in x and explain why there are no real solutions. [5]

- 5 The mass, m grams, of a radioactive substance, present at time t years after being observed, is given by the formula $m = 195(0.8)^t$.
 - (i) Find
 - (a) the initial mass of the substance, [1]
 - (b) the mass of the substance when $t = 6$, [1]
 - (c) the value of t when the mass of the substance is $\frac{1}{4}$ of its initial mass. [4]

Give your answer correct to three significant figures.
 - (ii) Explain why the mass of the substance can never be more than 195. [1]
 - (iii) Sketch the graph of m against t , where $t > 0$. [1]

Answer Key

1(a)	$2 + u$
1(b)	$u - 3$
1(c)	$\frac{3}{u}$
2(a)	$x = 2\frac{1}{4}$
2(b)	$y = 3\sqrt{3}, \frac{1}{3}$
3	$p = 0.378, 49$
4	$x^2 - 5x + 30 = 0$
5(i)(a)	195 g
5(i)(b)	51.1 g
5(i)(c)	6.21 years
5(ii)	$\text{As } t \rightarrow \infty$ $0.8^t \rightarrow 0$ $195(0.8^t) \rightarrow 0$
5(iii)	Sketch graph